arXiv:1301.5171v2 [math.AC] 28 Jan 2013

DEPTH OF SOME SPECIAL MONOMIAL IDEALS

DORIN POPESCU AND ANDREI ZAROJANU

Abstract. Let $I \supsetneq J$ be two square free monomial ideals of a polynomial algebra over a field generated in degree $\geq d$, resp. $\geq d+1$. When I contains precisely one generator of degree d, the other generators having degrees $\geq d+1$, if the Stanley depth of I/J is $\leq d+1$ then the usual depth of I/J is $\leq d+1$ too, that is the Stanley Conjecture holds in these cases.

Key words : Monomial Ideals, Depth, Stanley depth.
2000 Mathematics Subject Classification: Primary 13C15, Secondary 13F20, 13F55, 13P10.

Introduction

Let $S = K[x_1, \ldots, x_n]$ be the polynomial algebra in n variables over a field K and $I \supsetneq J$ two monomial ideals of S generated by square free monomials of degrees $\geq d$, resp. $\geq d+1$ for some $d \in \mathbf{N}$. Then $\operatorname{depth}_S I/J \geq d$ (see [1, Proposition 3.1], [2, Lemma 1.1]) and upper bounds are given by Stanley's Conjecture if it holds. Finding such upper bounds is the subject of several papers [2], [3], [6], [4]. We remind below the notion of Stanley depth.

Let $P_{I\setminus J}$ be the poset of all square free monomials of $I \setminus J$ with the order given by the divisibility. Let $\mathcal{P}$ be a partition of $P_{I\setminus J}$ in intervals $[u,v] = \{w \in P_{I\setminus J} : u|w, w|v\}$, let us say $P_{I\setminus J} = \cup_i [u_i, v_i]$, the union being disjoint. Define $\operatorname{sdepth} \mathcal{P} = \min_i \deg v_i$ and the so called *Stanley depth* of I/J given by $\operatorname{sdepth}_S I/J = \max_{\mathcal{P}} \operatorname{sdepth} \mathcal{P}$, where $\mathcal{P}$ runs in the set of all partitions of $P_{I\setminus J}$ (see [1], [7]). Stanley's Conjecture says that $\operatorname{sdepth}_S I/J \geq \operatorname{depth}_S I/J$.

Let r be the number of the square free monomials of degree d of I and B (resp. C) be the set of the square free monomials of degrees $d+1$ (resp. $d+2$) of $I \setminus J$. Set $s = |B|$, $q = |C|$. If either $s > r+q$, or $r > q$, or $s < 2r$ then $\operatorname{sdepth}_S I/J \leq d+1$ and if the Stanley Conjecture holds then any of these numerical conditions would imply $\operatorname{depth}_S I/J \leq d+1$, independently of the characteristic of K. In particular this was proved directly in [3] and [6].

Suppose that $r = 1$. If $d = 1$ we showed in [4, Theorem 1.10] that almost always the Stanley Conjecture holds. It is the purpose of this note to complete the result for $d \geq 1$ in the next form.

Theorem *Suppose that $I \subset S$ is minimally generated by a square free monomial f of degree d, and a set of square free monomials of degrees $\geq d+1$. Assume that* $\operatorname{sdepth}_S I/J = d+1$. *Then* $\operatorname{depth}_S I/J \leq d+1$.

The support from grant ID-PCE-2011-1023 of Romanian Ministry of Education, Research and Innovation is gratefully acknowledged.

1. PROOF OF THE THEOREM

We may assume that $s \leq q+1$, otherwise apply [3] and $C \subset (f, B)$ by [4, Lemma 1.6]. Let E be the set of all square free monomials a of degrees $d+1$, which are not in (f) . We may suppose that $n > 2$ and there exists a square free monomial $c = fx_{n-1}x_n \notin J$ after renumbering the variables x, otherwise we may apply [4, Lemma 1.5]. Thus $q > 0$. Let $Z_c \subset S$ be the set containing all the monomials of degree $d+1$ such that $z|c$ and $z \notin (f)$, that is $z \neq fx_{n-1}, fx_n$. Then $Z_c \cap J = \emptyset$.

Suppose that we may find c with $Z_c \setminus I = Z_c$, that is $\{fx_{n-1}, fx_n\}$ are all divisors of c from I of degree $d+1$. Set $I' = (B \setminus \{fx_{n-1}, fx_n\})$ and $J' = I' \cap J$. We have necessary $B \supsetneq \{fx_{n-1}, fx_n\}$ because otherwise $\operatorname{sdepth}_S I/J = d+2$. Thus $I' \neq J'$. Note that no $b \in B$ dividing c belongs to I' and so $c \notin (J + I')$. In the following exact sequence

$$0 \to I'/J' \to I/J \to I/(I'+J) \to 0$$

the last term has sdepth $\geq d+2$ since $[f, c]$ is the whole poset of $(f)/(f) \cap (I'+J)$ except some monomials of degrees $\geq d+2$. It has also depth $\geq d+2$ because $x_{n-1}x_n \notin ((J+I') : f)$. The first term has sdepth $\leq \operatorname{sdepth}_S I/J = d+1$ by [5, Lemma 2.2] and so it has depth $\leq d+1$ by [2, Theorem 4.3]. It follows $\operatorname{depth}_S I/J \leq d+1$. Next suppose that there exist no such c, that is any $c \in C$ is a multiple of some $a \in E$. In particular, $E \neq \emptyset$. We will use induction on $|E|$, the case $E = \emptyset$ follows from above.

Let $a \in E$ and $I_a = (f, E \setminus \{a\})$, $J_a = I_a \cap J$. In the following exact sequence

$$0 \to I_a/J_a \to I/J \to I/(I_a + J) \to 0$$

the last term has depth $\geq d+1$ because it is isomorphic with $(a)/(a) \cap (I_a + J)$. If $\operatorname{sdepth} I_a/J_a \leq d+1$ then the first term has depth $\leq d+1$ by the induction hypothesis, so by the Depth Lemma we get $\operatorname{depth} I/J \leq d+1$. Thus we may assume that $\operatorname{sdepth} I_a/J_a \geq d+2$, for any $a \in E$ and let $\mathcal{P}_a$ be a partition on I_a/J_a with sdepth $= d+2$.

Now *suppose that* $s = q+1$. Then there exists one c such that $\mathcal{P}_a$ contains no interval $[b, c]$ with either $b = f$, or $b \in (B \setminus \{a\})$. We have $c \in (a')$ where $a' \in E$. Of course we have $a' \neq a$, even more $c \notin (a)$ because otherwise we can add to $\mathcal{P}_a$ the interval $[a, c]$ and get a partition for I/J with sdepth $\geq d+2$. Consider $\mathcal{P}_{a'}$ and c' as above. We have $c \neq c'$ because otherwise we get $a|c$. In $\mathcal{P}_a$ there exists an interval $[f, fx_ix_j]$ and similarly there exists in $\mathcal{P}_{a'}$ an interval $[f, fx'_ix'_j]$ with $i, j, i', j' \in [n] \setminus \operatorname{supp}(f)$. Also we have in $\mathcal{P}_a$ for all $b \in B \setminus \{a, fx_i, fx_j\}$ an interval $[b, c_b]$. We define $h : (B \setminus \{a\}) \to (C \setminus \{c\})$ by $b \to c_b$ and $h(fx_i) = h(fx_j) = fx_ix_j$ and let $g : (C \setminus \{c\}) \to (B \setminus \{a\})$ defined by $c_b \to b$ and $g(fx_ix_j) = fx_i$. Similarly we define h', g' for $\mathcal{P}_{a'}$.

We want to build a partition $\mathcal{P}$ for I/J with sdepth $= d+2$. This will be false ending the proof. Set $a_0 = a$ and by the recurrence $c_p = h'(a_p)$, $a_{p+1} = g(c_p)$ for $p \geq 0$. The construction stops at step $p = e$ if one of the following cases appears

1) $a_e = a'$,

2) $c_e = c$,

3) $a_e = fx_{j'}$ after $fx_{i'}$ already appeared among a_p with $p < e$.

In the first case we set $c_e = c$ (which is in fact the second case), and we see that h gives a bijection between $\{a_1, \dots, a_e\}$ and $\{c_0, \dots, c_{e-1}\}$. But h gives also a bijection between $B \setminus \{a, fx_j, a_1, \dots, a_e\}$ and $C \setminus \{c, c_0, \dots, c_{e-1}\}$. If we are not in case 3), that is we have not both $fx_{i'}$ and $fx_{j'}$ between a_p, then the intervals $[a_p, c_p]$, $0 \leq p \leq e$ and the intervals $[g(\tilde{c}), \tilde{c}]$, $\tilde{c} \in C \setminus \{c, c_0, \dots, c_{e-1}\}$ give a partition $\mathcal{P}$ of I/J with sdepth $\geq d+2$ if we change $a_p = fx_i$ or may be $g(\tilde{c}) = fx_i$ by f.

It remains to assume the case 3). Now we set $a_e = fx_j$, we can do this because $fx_j \notin \mathrm{Im}(g)$. We continue the procedure in the same way, $c_p = h'(a_p)$, $a_{p+1} = g(c_p)$ for $p \geq e$. The construction stops now for $p = u$ when either $a_u = a'$, or $c_u = c$. In the first subcase we set $c_u = c$ and continue with $a_p = g'(c_{p-1})$, $c_p = h(a_p)$ for $p > u$. Now the procedure may stop only for $p = t$ with $c_t = c'$. In this situation we see that h' gives a bijection between $\{a_0, \dots, a_{u-1}, a_{u+1}, \dots, a_t\}$ and $\{c_0, \dots, c_{t-1}\}$. It follows that h' gives also a bijection between $B \setminus \{fx_{j'}, a_0, \dots, a_t\}$ and $C \setminus \{c_0, \dots, c_t\}$. So the intervals $[a_p, c_p]$, $0 \leq p \leq t$ and the intervals $[\tilde{a}, h'(\tilde{a})]$, $\tilde{a} \in B \setminus \{fx_{j'}, a_0, ..., a_t\}$ give a partition $\mathcal{P}$ of I/J with sdepth $\geq d+2$ if we change $a_k = fx_{i'}$ with f.

In the last subcase we set $a_{u+1} = a'$ and we continue with $c_p = h(a_p)$, $a_{p+1} = g'(c_p)$, for $p > u$. Now the procedure may stop for $p = t$ with $c_t = c'$. In this situation we see that h' is a bijection between $\{a_0, \dots, a_u, a_{u+2}, \dots, a_t\}$ and $\{c_0, \dots, c_{t-1}\}$. It follows that h' is also a bijection between $B \setminus \{fx_{j'}, a_0, \dots, a_t\}$ and $C \setminus \{c_0, \dots, c_t\}$. So the intervals $[a_p, c_p]$, $0 \leq p \leq t$ and the intervals $[\tilde{a}, h'(\tilde{a})]$, $\tilde{a} \in B \setminus \{fx_{j'}, a_0, ..., a_t\}$ give a partition $\mathcal{P}$ of I/J with sdepth $\geq d+2$ if we change $a_k = fx_{i'}$ with f.

Next *suppose that* $s < 1 + q$. When $d = 1$ this was done in [4] and similarly one can do it for $d > 1$. Here we give a shorter form. Set $I_n = (B \setminus \{fx_n\})$, $J_n = I_n \cap J$. In the following exact sequence

$$0 \to I_n/J_n \to I/J \to I/(I_n + J) \to 0$$

the last term has sdepth and depth $\geq d+1$ since $[f, fx_n]$ is an interval of $(f)/(f) \cap (I_n + J)$ and $x_n \notin ((J + I_n) : f)$. If the first term has sdepth $= d+1$ then by [2, Theorem 4.3] we get depth $= d+1$ and applying the Depth Lemma the conclusion follows. So we can assume that there exists a partition $\mathcal{P}_n$ of I_n/J_n with sdepth $= d+2$. $\mathcal{P}_n$ contains an interval $[b, fx_{n-1}x_n]$, for some $b \in E$ because otherwise adding $[f, fx_{n-1}x_n]$ we get a partition of I/J with sdepth $= d+2$.

$\mathcal{P}_a$ contains an interval $[f, fx_ix_j]$, for some $i, j \in [n] \setminus \mathrm{supp}\, f$. Let $\mathcal{P}_i$, be a partition with sdepth $= d+2$ similar to $\mathcal{P}_n$ (here $I_i = (B \setminus \{fx_i\})$). We have in $\mathcal{P}_a$ for all $b \in B \setminus \{a, fx_i, fx_j\}$ an interval $[b, c_b]$. Almost as above, we define $h : (B \setminus \{a\}) \to C$ by $b \to c_b$ and $h(fx_i) = h(fx_j) = fx_ix_j$ and $g : C \to (C \cup (B \setminus \{a\}))$ by $c_b \to b$, $g(fx_ix_j) = fx_i$ and $g(c) = c$ if c is not present in an interval $[b, c_b]$ with $\deg b \leq d+1$. Similarly we define h', g' for $\mathcal{P}_i$. From above we have $fx_ix_j \in \mathrm{Im}\, h'$.

We build a partition $\mathcal{P}'_a$ with sdepth $= d+2$ on I'_a/J'_a, where $I'_a = (B \setminus \{a\})$, $J'_a = I' \cap J$. Consider $a_0 = a$ and set $c_{p-1} = h'(a_{p-1})$, $a_p = g(c_{p-1})$, for $p \geq 1$. If we get a c_p such that $g(c_p) = c_p$ then by changing in $\mathcal{P}_a$ the intervals $[a_t, c_{t-1}]$, $t \in [p]$ with the intervals $[a_t, c_t]$, $t \in [p]$ we obtain a partition of I/J with sdepth $= d+2$, which is false. The construction stops when $c_p = fx_ix_j$. Then set $a_{p+1} = fx_i$ and $c_{p+1} = fx_ix_k$, for some k. Let $\mathcal{P}'_a$ be given by the intervals $[a_{i+1}, c_i]$, $i \in [p-1]$,

$[fx_j, c_p]$, $[a_{p+1}, c_{p+1}]$, $[b, h(b)]$, for all $b \in B \setminus \{a_1, a_2, ..., a_{p+1}, fx_i\}$ and other intervals starting with monomials of degree $\geq d+2$.

Now we give a partition with sdepth $= d+2$ for I_B/J_B, where $I_B = (B)$ and $J_B = I_B \cap J$. This leads to a contradiction, because replacing, let us say, the interval $[fx_n, fx_nx_k]$ with $[f, fx_nx_k]$ induces a partition of I/J with sdepth $= d+2$. As $s < 1+q$ it follows that there exists c which is not in $\operatorname{Im} h$. Then $c \in (a)$ for some $a \in E$ as we have seen. We consider $\mathcal{P}'_a$ and the corresponding i, j and maps h, g as above. Similarly we define h', g' for $\mathcal{P}_i$. Set $a_0 = fx_n$ and $c_{p-1} = h(a_{p-1})$, $a_p = g'(c_{p-1})$, for $p \geq 1$. The construction may stop at step e if either $a_e = a$, or $c_e \notin \operatorname{Im} h'$. In the first case $[a_i, c_i]$, for $i \in [e-1]$, $[a, c]$, $[b, h'(b)]$, for all $b \in B \setminus \{a_0, a_1, ..., a_e\}$ and other intervals starting with monomials of degree $\geq d+2$ give a partition of I_B/J_B with sdepth $= d+2$. In the second case, $[a_i, c_i]$, for $i \in [e]$, $[a, c]$, $[b, h'(b)]$, for all $b \in B \setminus \{a_0, a_1, ..., a_e\}$ and other intervals starting with monomials of degree $\geq d+2$ give a partition of I_B/J_B with sdepth $= d+2$.

References

[1] J. Herzog, M. Vladoiu, X. Zheng, *How to compute the Stanley depth of a monomial ideal,* J. Algebra, **322** (2009), 3151-3169.

[2] D. Popescu, *Depth of factors of square free monomial ideals*, to appear in Proceedings AMS, arXiv:AC/1110.1963.

[3] D. Popescu, *Upper bounds of depth of monomial ideals*, to appear in J. Commutative Alg., (2012),arXiv:AC/1206.3977.

[4] D. Popescu, A. Zarojanu, *Depth of some square free monomial ideals*, Bull. Math. Soc. Sci. Math. Roumanie, **56(104)**, 2013,117-124.

[5] A. Rauf, *Depth and Stanley depth of multigraded modules*, Comm. Algebra, **38** (2010),773-784.

[6] Y.H. Shen, *Lexsegment ideals of Hilbert depth 1*, (2012), arxiv:AC/1208.1822v1.

[7] R. P. Stanley, *Linear Diophantine equations and local cohomology*, Invent. Math. **68** (1982) 175-193.

Dorin Popescu, "Simion Stoilow" Institute of Mathematics of Romanian Academy, Research unit 5, P.O.Box 1-764, Bucharest 014700, Romania

E-mail address: `dorin.popescu@imar.ro`

Andrei Zarojanu, Faculty of Mathematics and Computer Sciences, University of Bucharest, Str. Academiei 14, Bucharest, Romania

E-mail address: `andrei_zarojanu@yahoo.com`